

Anorthoclase

(Na, K)AlSi₃O₈

©2001 Mineral Data Publishing, version 1.2

Crystal Data: Triclinic. *Point Group:* $\bar{1}$. Short prismatic crystals; also tabular, rhombic, flattened along [010], to 5 cm. *Twinning:* Baveno, Carlsbad, and Manebach laws; polysynthetic albite and pericline law twinning produce a grid pattern on {100}.

Physical Properties: *Cleavage:* Perfect on {001}, less perfect on {010}; partings on {100}, {110}, $\{\bar{1}10\}$, and $\{\bar{2}01\}$. *Fracture:* Uneven. *Tenacity:* Brittle. *Hardness* = 6
D(meas.) = 2.57–2.60 D(calc.) = 2.57

Optical Properties: Transparent. *Color:* Colorless, also white, pale creamy yellow, red, green. *Streak:* White. *Luster:* Vitreous, may be pearly on cleavages.

Optical Class: Biaxial (-). *Orientation:* $Z \wedge b \simeq 5^\circ$. *Dispersion:* $r > v$, weak.
 $\alpha = 1.524\text{--}1.526$ $\beta = 1.529\text{--}1.532$ $\gamma = 1.530\text{--}1.534$ $2V(\text{meas.}) = 42^\circ\text{--}52^\circ$

Cell Data: *Space Group:* $C\bar{1}$. $a = 8.287$ $b = 12.972$ $c = 7.156$ $\alpha = 91.05^\circ$ $\beta = 116.26^\circ$
 $\gamma = 90.15^\circ$ $Z = 4$

X-ray Powder Pattern: Grande Caldeira, Azores. (ICDD 9-478).
3.211 (100), 3.243 (90), 4.106 (16), 2.162 (16), 6.49 (14), 3.768 (14), 3.726 (14)

Chemistry:	(1)
SiO ₂	62.79
Al ₂ O ₃	22.12
Fe ₂ O ₃	0.36
FeO	0.41
CaO	3.76
Na ₂ O	7.35
K ₂ O	2.98
H ₂ O ⁺	0.19
H ₂ O ⁻	0.07
Total	100.03

(1) Mt. Erebus, Ross Island, Antarctica; corresponds to $(\text{Na}_{0.64}\text{Ca}_{0.18}\text{K}_{0.17})_{\Sigma=0.99}$
 $(\text{Al}_{0.98}\text{Fe}_{0.02}^{2+}\text{Fe}_{0.01}^{3+})_{\Sigma=1.01}(\text{Si}_{2.81}\text{Al}_{0.19})_{\Sigma=3.00}\text{O}_8$.

Mineral Group: Feldspar (alkali) group; intermediate between low sanidine and high albite.

Occurrence: In high-temperature sodic volcanic and hypabyssal rocks.

Association: Typically in a fine-grained groundmass or weathered out as loose crystals.

Distribution: Rather abundant worldwide. Some localities for well-characterized material include: on Pantelleria and Ustica Islands, Italy. At Larvik, Norway. From Berkum, North Rhine-Westphalia, Germany. On Grande Caldeira Island, Azores. At Ropp, Nigeria. On Mt. Kenya, Kenya. From Kilimanjaro, Tanzania. At Chilposan, near Minchon, North Korea. From Ogaya, Toyama Prefecture, and Madarajima, Saga Prefecture, Japan. At Kakanui, New Zealand. From Mt. Anakie and Mt. Franklin, Daylesford, Victoria, Australia. Large crystals from Mt. Erebus, Ross Island, Antarctica. At Boron, Kern Co., California, USA.

Name: From the Greek for *oblique* and *fracture*, descriptive of the cleavage.

References: (1) Dana, E.S. (1892) Dana's system of mineralogy, (6th edition), 324–325. (2) Deer, W.A., R.A. Howie, and J. Zussman (1963) Rock-forming minerals, v. 4, framework silicates, 6–93. (3) Phillips, W.R. and D.T. Griffen (1981) Optical mineralogy, 342–344. (4) Carmichael, I.S.E. and W.S. MacKenzie (1964) The lattice parameters of high-temperature triclinic sodic feldspars. *Mineral. Mag.*, 33, 949–962. (5) Harlow, G.E. (1982) The anorthoclase structures: the effects of temperature and composition. *Amer. Mineral.*, 67, 975–996.

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system or transmitted in any form or by any means, electronic, mechanical, photocopying, recording, or otherwise without the prior written permission of Mineral Data Publishing.