

Bultfonteinite**Ca₂SiO₂(OH, F)₄**

©2001 Mineral Data Publishing, version 1.2

Crystal Data: Triclinic. *Point Group:* $\bar{1}$. As radiating prismatic acicular crystals and radial spherules, to 2 cm. *Twinning:* Interpenetrating on {100} and {010}; polysynthetic.

Physical Properties: *Cleavage:* Good on {100} and {010}. *Fracture:* Conchoidal. Hardness = 4.5 D(meas.) = 2.73 D(calc.) = 2.74

Optical Properties: Transparent. *Color:* Pale pink to colorless. *Luster:* Vitreous. *Optical Class:* Biaxial (+). *Orientation:* $Z' \wedge c = 27^\circ\text{--}29^\circ$ on {010}; $Z' \wedge c = 46^\circ\text{--}48^\circ$ on {100}. *Dispersion:* $r > v$, barely perceptible. $\alpha = 1.587(2)$ $\beta = [1.590]$ $\gamma = 1.597(2)$ $2V(\text{meas.}) = 70^\circ$

Cell Data: *Space Group:* $P\bar{1}$. $a = 10.992(4)$ $b = 8.185(2)$ $c = 5.671(1)$ $\alpha = 93^\circ 57'$ $\beta = 90^\circ 19'$ $\gamma = 89^\circ 51'$ $Z = 4$

X-ray Powder Pattern: Bultfontein mine, Kimberley, South Africa. 1.93 (100), 8.12 (60), 2.92 (60), 2.88 (60), 2.037 (50), 4.06 (40), 3.50 (40)

Chemistry:

	(1)	(2)
SiO ₂	26.50	24.06
B ₂ O ₃		0.02
Al ₂ O ₃	0.72	1.37
CaO	54.20	53.59
Na ₂ O		0.04
F	8.81	7.90
H ₂ O ⁺		10.78
H ₂ O ⁻		0.75
H ₂ O	13.36	
CO ₂		4.65
P ₂ O ₅		0.02
-O = F ₂	3.71	3.33
Total	99.88	99.85

(1) Bultfontein mine, Kimberley, South Africa. (2) Fuka, Japan; corresponds to (Ca_{2.04}(Si_{0.96}Al_{0.06})_{Σ=1.02}O_{2.13}[(OH)_{2.87}F_{1.00}]_{Σ=3.87}).

Occurrence: In a large "horse" of diabase and shale fragments in a kimberlite pipe (Bultfontein mine, Kimberley, South Africa); in a contact zone in thermally metamorphosed limestone (Crestmore, California, USA).

Association: Calcite, apophyllite, natrolite (Bultfontein mine, Kimberley, South Africa); awillite, scawtite (Crestmore, California, USA); oyelite, scawtite, xonotlite (Fuka, Japan).

Distribution: In South Africa, in Cape Province, in the Bultfontein and Dutoitspan diamond mines, Kimberley; the Jagersfontein diamond mine, Orange River Colony; and the N'Chwaning and Wessels manganese mines, near Kuruman. In the USA, at Crestmore, Riverside Co., California. In the Hatrurim Formation, Israel. In Japan, from Fuka, near Bichu, and in the Mihara mine, Okayama Prefecture.

Name: For the Bultfontein mine, Kimberley, South Africa, where it was first discovered.

Type Material: Cambridge University, Cambridge; The Natural History Museum, London, England, 1928,78.

References: (1) Parry, J., A.F. Williams, and F.E. Wright (1932) On bultfonteinite, a new fluorine-bearing hydrous calcium silicate from South Africa. *Mineral. Mag.*, 23, 145–162. (2) (1933) *Amer. Mineral.*, 18, 32 (abs. ref. 1). (3) McIver, E.J. (1963) The structure of bultfonteinite, Ca₄Si₂O₁₀F₂H₆. *Acta Cryst.*, 16, 551–558. (4) Kusachi, I., C. Henmi, and K. Henmi (1984) An oyelite-bearing vein at Fuka, the town of Bichu, Okayama Prefecture, Japan. *J. Japan. Assoc. Mineral. Petrol. Econ. Geol.*, 79, 267–275.

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system or transmitted in any form or by any means, electronic, mechanical, photocopying, recording, or otherwise without the prior written permission of Mineral Data Publishing.