

**Gorceixite****BaAl<sub>3</sub>(PO<sub>4</sub>)(PO<sub>3</sub>OH)(OH)<sub>6</sub>**

©2001-2005 Mineral Data Publishing, version 1

**Crystal Data:** Monoclinic, pseudorhombohedral. *Point Group:* *m*. Rare rhombic crystals. May be radial fibrous in spheroidal concretions, to 1 cm; in microcrystalline grains and pebbles.

**Physical Properties:** *Fracture:* Porcelaneous [*sic*]. *Hardness* = 6 *D*(meas.) = 3.036–3.185 *D*(calc.) = 3.389

**Optical Properties:** Transparent. *Color:* Colorless, green, brown, may be mottled; colorless in thin section. *Luster:* Vitreous.

*Optical Class:* Uniaxial (+); anomalous, with low birefringence. *Orientation:* Length positive, parallel extinction.  $\omega = 1.610$   $\epsilon = 1.620$

**Cell Data:** *Space Group:* *Cm* (probable). *a* = 12.195–12.225 *b* = 7.040–7.056 *c* = 7.055–7.061  $\beta = 125.10^\circ$ – $125.30^\circ$  *Z* = 2

**X-ray Powder Pattern:** Big Fish River, Canada.

3.010 (100), 5.769 (93), 3.529 (56), 2.292 (39), 1.9197 (29), 2.232 (23), 1.7626 (20)

<b>Chemistry:</b>		(1)	(2)		(1)	(2)	(1)	(2)
P <sub>2</sub> O <sub>5</sub>	28.11	27.76	BaO	28.9	29.99	S	0.28	
Al <sub>2</sub> O <sub>3</sub>	30.28	29.92	H <sub>2</sub> O	[9.8]	12.33	Total	[100.0]	100.00
SrO	0.28		F	2.3				

(1) Sydney Basin, Australia; by electron microprobe and X-ray fluorescence, H<sub>2</sub>O by difference; corresponds to (Ba<sub>0.95</sub>Sr<sub>0.01</sub>)<sub>Σ=0.96</sub>Al<sub>3.02</sub>(PO<sub>4</sub>)<sub>2.01</sub>[(OH), F]<sub>5.83</sub>. (2) BaAl<sub>3</sub>(PO<sub>4</sub>)(PO<sub>3</sub>OH)(OH)<sub>6</sub>.

**Mineral Group:** Crandallite group.

**Occurrence:** Probably always a secondary mineral; in novaculites and other sedimentary rocks; in carbonatites; from hydrothermal argillic alteration zones; in topaz greisen; in the fine-grained fraction of soils; commonly as favas in diamantiferous river sands.

**Association:** Dawsonite, alumohydrocalcite, nordstrandite (Sydney Basin, Australia); lazulite, siderite, “limonite”, quartz (Big Fish River, Canada); diamond (river sands, may be coincidental).

**Distribution:** In Brazil, in Minas Gerais, along the Rio Abaeté, Rio Bagagem, and Rio Douradinho, near Curralinho and Dattas, Diamantina, and in the Araxá carbonatite, containing an estimated 20 million tons; in Goyaz, in the Rio Paranaíba and at Veríssimo; in São Paulo, in the Rio Canoas and at Patrocínio de Sapucahy. At Somabula, Zimbabwe. From the Mrima Hill carbonatite, Kenya. At Issineru, Guiana. In the Bonsa River, near Dompim, Gold Coast. From the Oiyi district, Sierra Leone. In the Buffels River, Spektakel Valley, west of Springbok, Namaqualand, South Africa. From the Big Fish River–Rapid Creek area, Yukon Territory, Canada. In the Zig Zag Mountains, about 10 km northeast of Hot Springs, Garland Co., Arkansas; at the Silver Mine area, Madison Co., Missouri; from Indian Mountain, Cherokee Co., Alabama, USA. In the Hospital quarry, Elgin, Morayshire, Scotland. At the Lengenbach quarry, Binntal, Valais, Switzerland. In Australia, from Glen Alice and elsewhere in the Sydney Basin, New South Wales, and in the Moculta phosphate quarry, northeast of Angaston, South Australia; from the Magnet and Adelaide mines, Dundas, Tasmania. A few other localities are known.

**Name:** To honor Henrique Gorceix (1842–1919), French mining geologist and mineralogist, Director, School of Mines, Ouro Preto, Brazil.

**Type Material:** n.d.

**References:** (1) Palache, C., H. Berman, and C. Frondel (1951) Dana’s system of mineralogy, (7th edition), v. II, 833. (2) Radoslovich, E.W. and P.G. Slade (1980) Pseudo-trigonal symmetry and the structure of gorceixite. *Neues Jahrb. Mineral., Monatsh.*, 157–170. (3) Radoslovich, E.W. (1982) Refinement of gorceixite structure in *Cm*. *Neues Jahrb. Mineral., Monatsh.*, 446–464. (4) Blanchard, F.N. (1989) New X-ray powder data for gorceixite, BaAl<sub>3</sub>(PO<sub>4</sub>)<sub>2</sub>(OH)<sub>5</sub>•H<sub>2</sub>O, an evaluation of *d*-spacings and intensities, pseudosymmetry and its influence on the figure of merit. *Powder Diffraction*, 4(4), 227–230.

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system or transmitted in any form or by any means, electronic, mechanical, photocopying, recording, or otherwise without the prior written permission of Mineral Data Publishing.