

Hydroxykenomicrolite

(□, Na, Sb³⁺)₂Ta₂O₆(OH)

Crystal Data: Cubic. *Point Group:* 4/m $\bar{3}$ 2/m. As cubo-octahedral crystals; granular, to 3 mm.

Physical Properties: *Fracture:* Uneven. *Tenacity:* Brittle. Hardness = ~5
VHN = 670-780 (100 g load). D(meas.) = 6.4-6.6 D(calc.) = 6.49 Fluoresces yellow-orange to orange under LW UV; weak yellowish cathodoluminescence.

Optical Properties: Transparent; may be opaque except in thinnest fragments.
Color: Colorless to gray, yellow-orange, black. *Luster:* Adamantine to vitreous.
Optical Class: Isotropic. $n = > 1.8$
R: (480) 13.8, (551) 13.6, (589) 13.6, (656) 13.0

Cell Data: *Space Group:* Fd $\bar{3}$ m. $a = 10.496\text{-}10.515$ Z = 8

X-ray Powder Pattern: Kola Peninsula, Russia.
3.04 (10), 1.860 (10), 1.587 (10), 1.012 (10), 3.17 (9), 1.370 (9), 1.017 (9)

| Chemistry: | (1) | (2) | (3) | | (1) | (2) | (3) |
|--------------------------------|------|------|------|-------------------|-------|--------|--------|
| Nb ₂ O ₅ | 2.3 | 2.8 | 1.2 | CaO | 0.1 | | 0.6 |
| Ta ₂ O ₅ | 72.0 | 70.8 | 72.5 | Na ₂ O | 1.3 | 1.7 | 2.4 |
| Bi ₂ O ₃ | 0.7 | 0.3 | 0.6 | K ₂ O | | 0.0 | 0.05 |
| Sb ₂ O ₃ | 13.6 | 14.2 | 9.7 | Cs ₂ O | 7.3 | 7.4 | 5.4 |
| SnO | | | 0.1 | H ₂ O | [1.5] | [1.2] | |
| PbO | 1.6 | 0.8 | 5.3 | Total | 98.9 | [99.5] | [99.0] |

(1) Kola Peninsula, Russia; by electron microprobe, average of five analyses; (OH)¹⁻ confirmed by IR spectroscopy. (2) Do.; by electron microprobe, H₂O calculated from structure; corresponds to [Cs_{0.31}(OH, F)_{0.69}]_{Σ=1.00}(Sb_{0.57}Na_{0.31}Pb_{0.02}Bi_{0.01})_{Σ=0.91}(Ta_{1.88}Nb_{0.12})_{Σ=2.00}[O_{5.69}(OH, F)_{0.31}]_{Σ=6.00}. (3) Tanco pegmatite, Canada; by electron microprobe, H₂O calculated from structural considerations; corresponds to [Cs_{0.22}K_{0.01}(OH, F)_{0.55}]_{Σ=0.78}(Na_{0.45}Sb_{0.40}Pb_{0.14}Ca_{0.06}Bi_{0.02}Sn_{0.01})_{Σ=1.08}(Ta_{1.95}Nb_{0.05})_{Σ=2.00}[O_{5.78}(OH, F)_{0.22}]_{Σ=6.00}.

Mineral Group: Pyrochlore supergroup (general formula - A₂B₂X₆Y); microlite group (B = Ta⁵⁺).

Occurrence: A rare accessory in some highly fractionated granite pegmatites.

Association: Simpsonite, stibiotantalite, microlite, wodginite, tantalite, pollucite, sosedkoite (Kola Peninsula, Russia); kimrobinsonite, manganotantalite, antimonian microlite (Mt. Holland, Western Australia).

Distribution: From the Leshaiia pegmatite, Vuorijarvi carbonatite complex, Kola Peninsula, Russia. In the Czech Republic, at the Dobrá Voda pegmatite, Velké Meziříčí. In the Tanco pegmatite, Bernic Lake, Manitoba, Canada. From near Mt. Holland, Western Australia.

Name: For a member of the *microlite* group with prefixes to indicate dominant OH⁻ (*hydroxy*) in the Y site and essential vacancies (*keno*) in the A site. Formerly ‘cesstibtantite’.

Type Material: Mining Museum, St. Petersburg (1324/1); Geology Museum, Kola Branch, Academy of Sciences, Apatity (5518); A.E. Fersman Mineralogical Museum, Academy of Sciences, Moscow, Russia (80827, 81058, vis6317).

References: (1) Voloshin, A.V., Y.P. Men'shikov, Y.A. Pakhomovskii, and L.I. Polezhaeva (1981) Cesstibtantite, (Cs, Na)SbTa₄O₁₂ - a new mineral from granitic pegmatites. Zap. Vses. Mineral. Obshch., 110, 345-351 (in Russian). (2) (1982) Amer. Mineral., 67, 413-414 (abs. ref. 1). (3) Atencio, D., M.B. Andrade, A.G. Christy, R. Gieré, and P.M. Kartashov (2010) The pyrochlore supergroup of minerals: nomenclature. Can. Mineral., 48, 673-698. (4) Nickel, E.H. and B.W. Robinson (1985) Kimrobinsonite, a new tantalum mineral from Western Australia, and its association with cesstibtantite. Can. Mineral., 23, 573-576.