


Janggunitite



©2001-2005 Mineral Data Publishing, version 1

Crystal Data: Orthorhombic. *Point Group:* n.d. As flaky crystals, to 0.13 mm, elongated || cleavage and flattened on {010}, probable; in radiating groups, dendritic or arborescent masses, and in colloform bands and very fine-grained aggregates.

Physical Properties: *Cleavage:* One direction, perfect. *Tenacity:* Very fragile [sic].
Hardness = 2–3 D(meas.) = 3.59 D(calc.) = 3.58

Optical Properties: Opaque. *Color:* Black; grayish white to gray in reflected light.
Streak: Brownish black to dark brown. *Luster:* Dull.
Optical Class: Biaxial. *Pleochroism:* Distinct; whitish to light gray in oil. *Anisotropism:* Very strong; yellowish brown with bluish tint, to gray. *Bireflectance:* Observed.
R₁–R₂: 13–15

Cell Data: *Space Group:* n.d. $a = 9.324$ $b = 14.05$ $c = 7.956$ $Z = 4$

X-ray Powder Pattern: Janggun mine, South Korea.
9.34 (s), 7.09 (s), 3.547 (s), 3.101 (s), 4.62 (m), 4.17 (m), 2.469 (m)

Chemistry:

	(1)
MnO ₂	74.91
Fe ₂ O ₃	4.19
MnO	11.33
PbO	0.03
H ₂ O ⁺	9.46
Total	99.92

(1) Janggun mine, South Korea; Fe and Pb by electron microprobe, total Fe as Fe₂O₃, H₂O as OH verified by IR; corresponds to Mn_{4.86}⁴⁺(Mn_{0.90}²⁺Fe_{0.30}³⁺)_{Σ=1.20}O_{8.09}(OH)_{5.92}.

Occurrence: Formed at a late stage of oxidation in a cementation zone of a manganese deposit.

Association: Nsutite, todorokite, calcite, rhodochrosite.

Distribution: In the Janggun mine, Bonghwa district, South Korea.

Name: For the Janggun mine, South Korea, its first locality.

Type Material: National Museum of Natural History, Washington, D.C., USA, 143150.

References: (1) Kim, S.J. (1977) Janggunitite, a new manganese hydroxide mineral from the Janggun mine, Bonghwa, Korea. *Mineral. Mag.*, 41, 519–523. (2) (1978) *Amer. Mineral.*, 63, 794 (abs. ref. 1).