

Crystal Data: Hexagonal. *Point Group:* 6/*m*. Crystals are typically hexagonal prismatic {11 $\bar{2}$ 0}, may be tapered by vincinal forms, with pyramidal {10 $\bar{1}$ 1} termination, to 10 cm.

Physical Properties: *Fracture:* Conchoidal. Hardness = 6.5–7.5 D(meas.) = 3.29(1) D(calc.) = 3.284 [F¹⁻ end-member]; 3.198 [(OH)¹⁻ end-member]. Piezoelectric.

Optical Properties: Transparent. *Color:* Colorless, pale yellowish brown, pale to dark cornflower-blue; colorless to pale blue in transmitted light, typically sector zoned.

Luster: Vitreous.

Optical Class: Uniaxial (–), may be anomalously biaxial (–) in sector-zoned crystals.

Pleochroism: Colorless to pale blue-violet if colored. *Dispersion:* $r > v$, distinct.

$\omega = 1.644$ – 1.647 $\epsilon = 1.637$ $\alpha = 1.637$ $\beta = 1.644$ – 1.647 $\gamma = 1.644$ – 1.652

2V(meas.) = 18°–33°

Cell Data: *Space Group:* P6₃/*m*. $a = 8.5591(3)$ $c = 8.1814(6)$ $Z = 2$

X-ray Powder Pattern: Mt. Sektui, Russia.

4.282 (100), 3.791 (57), 2.055 (44), 2.196 (35), 2.650 (33), 2.470 (30), 1.391 (30)

Chemistry:

	(1)	(2)
SiO ₂	0.03	0.17
B ₂ O ₃	33.51	33.81
Al ₂ O ₃	59.90	59.79
Fe ₂ O ₃	0.01	0.14
F	10.18	10.23
H ₂ O	n.d.	n.d.
–O = F ₂	4.29	4.31
Total	99.34	99.83

(1) Mt. Sektui, Russia; by electron microprobe. (2) Cape Cross, Namibia; by electron microprobe, (OH)¹⁻ calculated for charge balance; corresponds to Al_{6.00}Fe_{0.01}(B_{0.99}O₃)₅[F_{2.76}(OH)_{0.24}]_{Σ=3.00}.

Occurrence: A rare late hydrothermal mineral formed in granitic pegmatites.

Association: Albite, tourmaline, quartz, gypsum (Cape Cross, Namibia).

Distribution: Large crystals from Mt. Sektui, Nerschinsk district, Adun-Chilon Mountains, Siberia, Russia. In the Kukurt massif, Pamir Mountains, Tajikistan. In Namibia, gem crystals from a prospect near Cape Cross, at Mile 72, about 100 km north of Swakopmund, and from Ameib, Erongo Mountains. At Wannerköpfe and Emmelberg, near Üdersdorf, Eifel district, Germany.

Name: Honors Pavel Vladimirovich Eremeev (Jeremejev, *German*) (1830–1899), Russian mineralogist and crystallographer, who first recognized this species.

Type Material: Mining Institute, St. Petersburg, Russia, 412/1.

References: (1) Palache, C., H. Berman, and C. Frondel (1951) Dana's system of mineralogy, (7th edition), v. II, 330–332. (2) Herting, S. and H. Strunz (1978) Jeremejewit von Cape Cross in SW-Afrika. Aufschluss, 29, 45–53 (in German). (3) Foord, E.E., R.C. Erd, and G.R. Hunt (1981) New data for jeremejevite. Can. Mineral., 19, 303–310. (4) Sokolova, E.V., Y.K. Yegorov-Tismenko, S.V. Kargal'tsev, V.A. Klyakhin, and V.S. Urusov (1987) Refinement of the crystalline structure of synthetic fluorous jeremejevite Al₆[BO₃]₅F₃. Moscow Univ. Geol. Bull., 42, 79–81. (5) Pekov, I.V. (1998) Minerals first discovered on the territory of the former Soviet Union. Ocean Pictures, Moscow, 106–107.

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system or transmitted in any form or by any means, electronic, mechanical, photocopying, recording, or otherwise without the prior written permission of Mineral Data Publishing.