


Katophorite

(c)2001 Mineral Data Publishing, version 1.2

Crystal Data: Monoclinic. *Point Group:* 2/m. Prismatic, granular, also as fibrous aggregates or dendrites and skeletal crystals. Commonly rimming other minerals.

Twinning: || {100}.

Physical Properties: *Cleavage:* Perfect on {110}, intersecting at ~56° and ~124°; parting on {010}. *Tenacity:* Brittle. Hardness = 5–6 D(meas.) = 3.2–3.5 D(calc.) = [3.31]

Optical Properties: Transparent to translucent. *Color:* Black, dark green-black, bluish black; reddish yellow, bluish green in thin section. *Luster:* Vitreous.

Optical Class: Biaxial (-). *Pleochroism:* Strong; reddish yellow, reddish brown, and dark green to black. *Orientation:* Z = b. *Dispersion:* r > v. *Absorption:* Z > Y > X. $\alpha = 1.639\text{--}1.681$
 $\beta = 1.658\text{--}1.688$ $\gamma = 1.600\text{--}1.690$ 2V(meas.) = ~0°–50°

Cell Data: *Space Group:* C2/m. a = 10.019(2) b = 18.036(7) c = 5.286(3)
 $\beta = 104.98(3)^\circ$ Z = 2

X-ray Powder Pattern: n.d.

Chemistry:	(1)	(2)	(1)	(2)
SiO ₂	48.04	45.98	MnO	1.11
TiO ₂	2.09	1.92	MgO	6.42
Al ₂ O ₃	3.86	2.79	CaO	8.08
Cr ₂ O ₃	0.09		Na ₂ O	4.18
FeO	25.58	28.97	K ₂ O	1.34
			Total	100.79
				97.83

(1) Baie-des-Moutons complex, Canada; by electron microprobe, corresponds to $(\text{Na}_{0.54}\text{K}_{0.26})_{\Sigma=0.80}(\text{Ca}_{1.31}\text{Na}_{0.69})_{\Sigma=2.00}(\text{Fe}_{3.25}^{2+}\text{Mg}_{1.45}\text{Ti}_{0.23}\text{Mn}_{0.15})_{\Sigma=5.08}(\text{Si}_{7.29}\text{Al}_{0.69}\text{Ti}_{0.01}\text{Cr}_{0.01})_{\Sigma=8.00}\text{O}_{22}(\text{OH})_2$. (2) Rallier-du-Baty Peninsula, Kerguelen Island; by electron microprobe, corresponds to $(\text{Na}_{0.80}\text{K}_{0.25})_{\Sigma=1.05}(\text{Ca}_{0.98}\text{Na}_{0.70}\text{Fe}_{0.32}^{2+})_{\Sigma=2.00}(\text{Fe}_{3.54}^{2+}\text{Mg}_{1.24}\text{Mn}_{0.15}\text{Ti}_{0.07})_{\Sigma=5.00}(\text{Si}_{7.32}\text{Al}_{0.52}\text{Ti}_{0.16})_{\Sigma=8.00}\text{O}_{22}(\text{OH})_2$.

Polymorphism & Series: Forms a series with magnesiokatophorite.

Mineral Group: Amphibole (sodic-calcic) group: $\text{Mg}/(\text{Mg} + \text{Fe}^{2+}) < 0.5$; $(\text{Na} + \text{K})_{\text{A}} \geq 0.5$; $0.67 \text{ Na}_{\text{B}} \text{ 1.33; } (\text{Ca} + \text{Na})_{\text{B}} \geq 1.34$; 6.5 Si 7.49.

Occurrence: In alkalic volcanic and plutonic igneous rocks; in blueschist facies jadeitites.

Association: Arfvedsonite, aegirine, nepheline, pyroxenes, eckermannite, chromite.

Distribution: May occur in the Oslo (Christiania) district, Norway, from where it was originally described. On the Rallier-du-Baty Peninsula, Kerguelen Island, in the south Indian Ocean. In the Baie-des-Moutons complex, La Tabatière, Quebec, Canada. At Tawmaw, Kachin State, northern Myanmar (Burma).

Name: From the Greek for *a carrying down*, in allusion to its volcanic origin.

Type Material: n.d.

References: (1) Dana, E.S. (1899) Dana's system of mineralogy, (6th edition), app. I, 14 [catophorite]. (2) Deer, W.A., R.A. Howie, and J. Zussman (1963) Rock-forming minerals, v. 2, chain silicates, 359–363. (3) Giret, A., B. Bonin, and J.-M. Leger (1980) Amphibole compositional trends in oversaturated and undersaturated alkaline plutonic ring complexes. *Can. Mineral.*, 18, 481–495. (4) Lalonde, A.E. and R.F. Martin (1983) The Baie-des-Moutons syenitic complex, La Tabatière, Québec, II. The ferromagnesian minerals. *Can. Mineral.*, 21, 81–91. (5) Hawthorne, F.C. (1983) The crystal chemistry of the amphiboles. *Can. Mineral.*, 21, 173–480, esp. 363. (6) Mével, C. and J.-R. Kiénaast (1986) Jadeite-kosmochlor solid solution and chromian sodic amphiboles in jadeitites and associated rocks from Tawmaw (Burma), *Bull. Minéral.*, 109, 617–633. (7) Phillips, W.R. and D.T. Griffen (1981) Optical mineralogy, 241–242.

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system or transmitted in any form or by any means, electronic, mechanical, photocopying, recording, or otherwise without the prior written permission of Mineral Data Publishing.