

Crystal Data: Cubic; may be metamict. *Point Group:* $4/m\bar{3}2/m$. Rare rounded octahedra with small cube modifications, to 8 mm; commonly anhedral, granular, in veinlets and aggregates.

Physical Properties: *Fracture:* [Conchoidal to uneven] (by analogy to pyrochlore group). *Tenacity:* Brittle. Hardness = ~5 D(meas.) = 5.30 (on impure material). D(calc.) = 5.19

Optical Properties: Semitransparent. *Color:* Dark brown to brown-black; brown in transmitted light; medium gray in reflected light. *Streak:* Pale brown to cream. *Luster:* Vitreous. *Optical Class:* Isotropic. $n = > 1.78$

Cell Data: *Space Group:* $Fd\bar{3}m$. $a = 10.351$ $Z = 8$

X-ray Powder Pattern: Věžná, Czech Republic; after heating at 700° C for one hour. 2.988 (10), 1.829 (6), 1.561 (5), 2.581 (4), 5.94 (3), 1.495 (3), 1.187 (3)

Chemistry:	(1)	(1)
Nb ₂ O ₅	21.6	PbO 0.13
Ta ₂ O ₅	19.3	SnO 2.9
TiO ₂	16.5	CaO 14.5
Al ₂ O ₃	0.49	Na ₂ O 0.30
Sb ₂ O ₃	23.2	F 0.15
FeO	0.6	H ₂ O 0.44
MnO	0.6	<u>-O = F₂ 0.06</u>
		Total 100.65

(1) Věžná, Czech Republic; by electron and ion microprobe; total Sb as Sb₂O₃, Fe as FeO, Mn as MnO, Sn as SnO; corresponds to (Ca_{1.11}Sb³⁺_{0.69}Sn_{0.09}Fe_{0.04}Mn_{0.04}Na_{0.04})_{Σ=2.01}(Ti_{0.89}Nb_{0.70}Ta_{0.38}Al_{0.04})_{Σ=2.01}O₆[O_{0.76}(OH)_{0.21}F_{0.03}]_{Σ=1.00}.

Mineral Group: Pyrochlore supergroup (general formula - A₂B₂X₆Y); pyrochlore group (B = Nb⁵⁺).

Occurrence: As replacement masses and in veinlets, in a granite pegmatite in a serpentinite.

Association: Columbite, niobian rutile, antimony, stokesite, cassiterite, zircon, albite.

Distribution: From Věžná, Czech Republic.

Name: For a member of the *pyrochlore* group with prefixes to indicate dominant oxygen (*oxy*) in the Y site and dominant calcium (*calcio*) in the A site. Formerly 'stibiobetafite'.

Type Material: University of Manitoba, Winnipeg (M5233); Royal Ontario Museum, Toronto, Canada (M35630).

References: (1) Černý, P., F.C. Hawthorne, J.H.G. Laflamme, and J.R. Hinthorne (1979) Stibiobetafite, a new member of the pyrochlore group from Věžná, Czechoslovakia. *Can. Mineral.*, 17, 583-588. (2) (1981) *Amer. Mineral.*, 66, 1278 (abs. ref. 1). (3) Atencio, D., M.B. Andrade, A.G. Christy, R. Gieré, and P.M. Kartashov (2010) The pyrochlore supergroup of minerals: nomenclature. *Can. Mineral.*, 48, 673-698.